

13:30 – 14:30	Room: Anadolu Auditorium	FC6
Session type:	Free Communication Session 6	
Session title:	Diabetes and Insulin Resistance	
Moderators:	Claire Levy-Marchal, Paris, France Hülya Günöz, İstanbul, Turkey	

Aortic intima media thickness is a more sensitive marker of early atherosclerosis than carotid intima media thickness in children with type I diabetes FC6-98

Jennifer Harrington¹; Alexia Pena¹; Roger Gent²; Lino Piotto²; Craig Hirte³; Jennifer Couper⁴

¹Adelaide Women's and Children's Hospital, Endocrine and Diabetes, Adelaide, Australia; ²Adelaide Women's and Children's Hospital, Radiology, Adelaide, Australia; ³Adelaide Women's and Children's Hospital, public health research unit, Adelaide, Australia; ⁴University of Adelaide, Paediatrics, Adelaide, Australia

Insulin, C-peptide concentrations and insulin sensitivity in children with Cystic Fibrosis related diabetes FC6-99

Stephen Oriordan¹; Peter Hindmarsh¹; Costigan Colm²; Nuala Murphy³; Edna Roche⁴; Hilary Hoey⁴

¹The Institute of Child Health, Developmental Endocrinology Research Group, London, United Kingdom; ²Our Lady's Hospital Crumlin, Paediatric Diabetes and Endocrinology, Dublin, Ireland; ³The Children's University Hospital, Paediatric Diabetes and Endocrinology, Dublin, Ireland; ⁴The National Childrens Hospital Tallaght, Paediatric Diabetes and Endocrinology, Dublin, Ireland

Prospective, placebo-controlled, randomized treatment of 67 obese children/adolescents with metformin FC6-100

Susanna Wiegand¹; Dagmar L'Allemand²; Hanna Hübel¹; Mareike Bürmann¹; Reinhard Holl³; Heiko Krude¹; Annette Grüters¹

¹Charité; Universitätsmedizin Berlin, Pediatric Endocrinology and Diabetology, Berlin, Germany; ²Ostschweizer Kinderspital, Pediatric Endocrinology and Diabetology, St. Gallen, Switzerland; ³University of Ulm, Epidemiology, Ulm, Germany

Insulin secretion and resistance in severely obese children due to Melanocortin-4 receptor gene mutation FC6-101

David Zangen¹; Ziva Ben Neriah²; Rafael Neshet³; Maha Abdulhadi-Atwan¹

¹Hadassah Hebrew University Medical Center, Division of Pediatric Endocrinology, Jerusalem, Israel; ²Hadassah Hebrew University Medical Center, Department of Genetics, Jerusalem, Israel; ³Hadassah Hebrew University Medical Center, Endocrine Service, Department of Internal Medicine, Jerusalem, Israel

Variations in the melanocortin 4 receptor gene is associated with difficulties to maintain this weight loss after lifestyle intervention in obese children FC6-102

Thomas Reinehr¹; Anke Hinney²; Andre Michael Toschke³; Johannes Hebebrand⁴

¹Vestische Youth Hospital, University of Witten Herdecke, Datteln, Germany; ²University of Duisburg-Essen, Pediatric Psychiatry, Essen, Germany; ³King's College London, Division of Health and Social Care Research, London, United Kingdom; ⁴University of Essen-Duisburg, Pediatric Psychiatry, Essen, Germany

A common functional variant in the p110 β gene promoter acts as a cis-acting eQTL (expression quantitative trait locus) to protect obese adolescents from insulin resistance FC6-103

C Le Stunff¹; A Dechartres¹; E Miraglia Del Giudice²; D Meyre³; P Bougneres⁴

¹St-Vincent de Paul hospital, INSERM U561, Paris, France; ²Second University of Naples, Department of Pediatrics, Napoli, Italy; ³Pasteur Institute, Lille, France; ⁴St-Vincent de Paul hospital, Pediatric Endocrinology and INSERM U561, Paris, France